arXiv:0809.4817v1 [gr-qc] 28 Sep 2008

EMP AND LINEAR SCHRÖDINGER MODELS FOR A CONFORMALLY BIANCHI I COSMOLOGY

Jennie D'Ambroise
Department of Mathematics and Statistics
University of Massachusetts
Amherst, MA 01003, USA
e-mail: dambroise@math.umass.edu

Abstract: Using a conformal version of the Bianchi I metric and a perfect fluid energy-momentum tensor, we show that the resulting Einstein field equations are equivalent to a generalized Ermakov-Milne-Pinney equation. Using a transformation analogous to the known correspondence between the classical EMP and a linear Schrödinger equation, we then construct the linear Schrödinger formulation for this cosmological model. Also, a note is included to show the extra term that arises when the energy-momentum tensor is taken to include a second term in addition to the scalar field.

AMS Subj. Classification: 83C05, 83C15
Key Words: Einstein field equations, Bianchi I universe, perfect fluid, Ermakov-Milne-Pinney (EMP), Schrödinger equation.

1. Introduction

A number of recent papers have explicated the Ermakov-Milne-Pinney (EMP) and Schrödinger content of various scalar field cosmologies, including both 2+1 and 3+1 curved FRLW [4, 5, 6, 7] and some anisotropic Bianchi models[1, 3, 8]. These connections are interesting since such equations often appear in condensed matter physics as well as quantum field theory; that is, such reformulations open the door to directly relating gravitational and nongravitational systems. For the conformally Bianchi I metric that we will consider here, F. L. Williams [8] has explicated a transformation which reduces the system of Einstein's field equations to a generalized EMP equation whose coefficient functions are coupled to a second equation. Here we show a similar transformation which results in an uncoupled generalized EMP equation. By implementing a transformation analogous to that which is known to relate the classical EMP to a linear Schrödinger equation, we further reduce the field equations to a linear Schrödinger equation. The author thanks Floyd L. Williams for his support and advice on the results presented here.

2. Einstein Equations

Consider the Einstein field equations $G_{ij} = K^2 T_{ij}$ for metric

$$ds^2 = -[a(t)b(t)c(t)]^2 dt^2 + a(t)^2 dx^2 + b(t)^2 dy^2 + c(t)^2 dz^2 \tag{2.1}$$

where $a(t), b(t), c(t) > 0$. The perfect fluid energy-momentum tensor is taken to be $T_{ij} = -\phi_{;i}\phi_{;j} + g_{ij}\left[\frac{1}{2}g^{k\lambda}\phi_{;k}\phi_{;\lambda} + V \circ \phi\right]$ for some scalar field ϕ with potential V. Under these assumptions the field equations become

$$\begin{aligned}
\frac{\dot{a}\dot{b}}{ab}+\frac{\dot{a}\dot{c}}{ac}+\frac{\dot{b}\dot{c}}{bc} &\overset{(i)}{=} K^2\left[\frac{\dot{\phi}^2}{2}+(abc)^2\left(V\circ\phi\right)\right] \\
\frac{\ddot{b}}{b}-\frac{\dot{b}^2}{b^2}+\frac{\ddot{c}}{c}-\frac{\dot{c}^2}{c^2}-\frac{\dot{a}\dot{b}}{ab}-\frac{\dot{a}\dot{c}}{ac}-\frac{\dot{b}\dot{c}}{bc} &\overset{(ii)}{=} K^2\left[-\frac{\dot{\phi}^2}{2}+(abc)^2\left(V\circ\phi\right)\right] \\
\frac{\ddot{a}}{a}-\frac{\dot{a}^2}{a^2}+\frac{\ddot{c}}{c}-\frac{\dot{c}^2}{c^2}-\frac{\dot{a}\dot{b}}{ab}-\frac{\dot{a}\dot{c}}{ac}-\frac{\dot{b}\dot{c}}{bc} &\overset{(iii)}{=} K^2\left[-\frac{\dot{\phi}^2}{2}+(abc)^2\left(V\circ\phi\right)\right] \\
\frac{\ddot{a}}{a}-\frac{\dot{a}^2}{a^2}+\frac{\ddot{b}}{b}-\frac{\dot{b}^2}{b^2}-\frac{\dot{a}\dot{b}}{ab}-\frac{\dot{a}\dot{c}}{ac}-\frac{\dot{b}\dot{c}}{bc} &\overset{(iv)}{=} K^2\left[-\frac{\dot{\phi}^2}{2}+(abc)^2\left(V\circ\phi\right)\right]
\end{aligned} \tag{2.2}$$

where $K^2 = 8\pi G$ and G is Newton's constant.

3. EMP Formulation

As in [8], equating pairs of the left sides of (ii)-(iv) in (2.2), allows us to deduce that

$$b(t) = C_1 e^{\lambda t} a(t), \qquad c(t) = C_2 e^{\mu t} a(t) \tag{3.1}$$

for real numbers $C_1, C_2 > 0$ and λ, μ. Forming the combination $3(i) - (ii) - (iii) - (iv)$ and using the relations (3.1), we obtain

$$4\frac{\dot{a}^2}{a^2} - \frac{\ddot{a}}{a} + 2(\lambda+\mu)\frac{\dot{a}}{a} + \lambda\mu = \frac{K^2}{2}\dot{\phi}^2. \tag{3.2}$$

Introducing the quantity $\beta(t) = e^{(\lambda+\mu)t} a(t)^3$, equation (3.2) becomes

$$2\frac{\dot{\beta}^2}{\beta^2} - \frac{\ddot{\beta}}{\beta} - (\lambda^2 - \lambda\mu + \mu^2) = \frac{3}{2}K^2\dot{\phi}^2. \tag{3.3}$$

Now find some function $\overline{\tau}(t)$ such that

$$\dot{\overline{\tau}}(t) = \theta\beta(t)^{(6+n)/6} \tag{3.4}$$

for any $n \neq 0$, $\theta > 0$ and define $\phi_1(\tau) = \phi(\overline{f}(\tau))$ where $\overline{f}(\tau)$ is the inverse of $\overline{\tau}(t)$. Then by (3.3), one can show that

$$y(\tau) = \beta(\overline{f}(\tau))^{n/6} \qquad \text{and} \qquad Q(\tau) = \frac{nK^2}{4}\phi_1'(\tau)^2 \tag{3.5}$$

satisfy the generalized EMP

$$y''(\tau) + Q(\tau)y(\tau) = \frac{-n(\lambda^2 - \lambda\mu + \mu^2)}{6\theta^2 y(\tau)^{(12+n)/n}}. \tag{3.6}$$

Similarly we may state the converse implication, which produces a solution of the Einstein equations (2.2) from a solution of the generalized EMP (3.6).

Theorem 1. *Suppose that $y(\tau)$ is a solution of the generalized EMP (3.6) for a known function $Q(\tau)$ and for some constants $n \neq 0$, $\theta > 0$ and λ, μ. Then a solution (a, b, c, ϕ, V) of (2.2) can be constructed as follows. First find functions $\overline{\tau}(t)$ and $\phi_1(\tau)$ such that*

$$\dot{\overline{\tau}}(t) = \theta y(\overline{\tau}(t))^{(6+n)/n}, \quad \phi_1'(\tau)^2 = \frac{4}{nK^2} Q(\tau). \tag{3.7}$$

Then the following solve the Einstein equations (i)-(iv) in (2.2):

$$a(t) = y(\overline{\tau}(t))^{2/n} e^{-\frac{1}{3}(\lambda+\mu)t}, \tag{3.8}$$

$$b(t) = C_1 e^{\lambda t} a(t), \quad c(t) = C_2 e^{\mu t} a(t), \quad \phi(t) = \phi_1(\overline{\tau}(t)), \tag{3.9}$$

$$V \circ \phi = \frac{\theta^2}{nK^2 C_1 C_2} \left[\frac{12}{n}(y')^2 - \frac{n(\lambda^2 - \lambda\mu + \mu^2)}{3\theta^2 (y)^{12/n}} - 2(y)^2 Q \right] \circ \overline{\tau} \tag{3.10}$$

for any constants $C_1, C_2 > 0$.

In the case $n = 6$, (3.6) reduces to the classical EMP equation, which is known to be equivalent to a linear Schrödinger equation. A similar transformation relates the generalized EMP in (3.6) to the linear Schrödinger equation

$$u''(x) + [E - P(x)]u(x) = 0. \tag{3.11}$$

This can be shown by letting $u(x) = y(\tau(g(x)))^{-6/n}$ where $g(x)$ is the inverse of $\sigma(t)$ which satisfies $\dot{\sigma}(t) = 1/\dot{\tau}(t)$, and where $\tau(t)$ is such that $\dot{\tau}(t) = \theta^{1/2} y(\tau(t))^{(6+n)/2n}$. Then $u(x)$ satisfies (3.11) for $E = -(\lambda^2 - \lambda\mu + \mu^2)$ and $P(x) = \frac{6\theta^2}{n} Q(\tau(g(x))) y(\tau(g(x)))^{(12+2n)/n}$. This motivates the next section, in which we directly relate Einstein's field equations (2.2) to the linear Schrödinger equation (3.11).

4. Linear Schrödinger Formulation

Beginning with a solution quintet (a, b, c, ϕ, V) of the Einstein equations (i)-(iv) in (2.2), one can follow the steps outlined above and obtain the formulas (3.1) for $b(t)$ and $c(t)$ in terms of $a(t)$ for some constants $C_1, C_2 > 0$ and λ, μ. Again, we deduce (3.2) and (3.3) for $\beta(t) = e^{(\lambda+\mu)t} a(t)^3$, find $\overline{\tau}(t)$ solving (3.4) for any $n \neq 0$ and $\theta > 0$, and define $\phi_1(\tau) = \phi(\overline{f}(\tau))$ where $\overline{f}(\tau)$ is the inverse of $\overline{\tau}(t)$. Now to obtain the Schrödinger formulation, we additionally solve for both $\tau(t)$ and $\sigma(t)$ that satisfy

$$\dot{\tau}(t) = \dot{\overline{\tau}}(\overline{f}(\tau(t)))^{1/2}, \qquad \dot{\sigma}(t) = \frac{1}{\dot{\tau}(t)}, \tag{4.1}$$

and we denote by $g(x)$ the inverse function of $\sigma(t)$. Introducing the quantities

$$u(x) = \beta(\overline{f}(\tau(g(x)))^{-1} \qquad \text{and} \qquad \psi(x) = \phi(\overline{f}(\tau(g(x)))), \tag{4.2}$$

one can show that $u(x)$, $P(x) = \frac{3K^2}{2}\psi'(x)^2$ and $E = -(\lambda^2 - \lambda\mu + \mu^2)$ solve the linear Schrödinger equation (3.11). We may now state the converse implication.

Theorem 2. *Suppose that $u(x)$ is a solution of the linear Schrödinger equation (3.11) for some known function $P(x)$ and constant $E < 0$. First find functions $\sigma(t)$ and $\tau(t)$ such that*

$$\dot{\sigma}(t) = u(\sigma(t))^{(6+n)/12} \quad \text{and} \quad \dot{\tau}(t) = \frac{1}{\dot{\sigma}(t)} \tag{4.3}$$

for some $n \neq 0$, and let $f \equiv \tau^{-1}$. Next find $\psi(x)$ such that $\psi'(x)^2 = \frac{2}{3K^2}P(x)$ and let λ, μ be any two constants such that

$$-(\lambda^2 - \lambda\mu + \mu^2) = E. \tag{4.4}$$

Finally, solve for $\overline{\tau}(t)$ such that

$$\dot{\overline{\tau}}(t) = \dot{\tau}(f(\overline{\tau}(t)))^2. \tag{4.5}$$

Then the following functions solve the Einstein equations (i)-(iv) in (2.2):

$$a(t) = \left[e^{(\lambda+\mu)t} u(\sigma(f(\overline{\tau}(t))))\right]^{-1/3}, \tag{4.6}$$

$$b(t) = C_1 e^{\lambda t} a(t), \quad c(t) = C_2 e^{\mu t} a(t), \quad \phi(t) = \psi(\sigma(f(\overline{\tau}(t)))), \tag{4.7}$$

$$V \circ \phi = \frac{1}{3K^2 C_1^2 C_2^2} \left[(u')^2 + [E - P]u^2\right] \circ \sigma \circ f \circ \overline{\tau}. \tag{4.8}$$

We note that including a second term in the energy-momentum tensor T_{ij} that has density $\rho(t)$ and pressure $p(t)$, one obtains a corresponding quantity

$$-\frac{nK^2}{4\theta^2} C_1^2 C_2^2 \frac{(\rho(\overline{f}(\tau)) + p(\overline{f}(\tau)))}{y(\tau)} \tag{4.9}$$

on the right side of the generalized EMP (3.6). Similarly (3.11) becomes nonlinear with the term

$$\frac{3K^2 C_1^2 C_2^2 (\overline{\rho}(x) + \overline{p}(x))}{2u(x)} \tag{4.10}$$

on the right side of the equation where $\overline{\rho}(x) \equiv \rho(\overline{f}(\tau(g(x))))$ and $\overline{p}(x) \equiv p(\overline{f}(\tau(g(x))))$. In this case both Theorems 1 and 2 include an additional $-\rho(t)$ added to the definitions of $V \circ \phi$.

5. Examples

Take the solution $u(x) = Ae^{-\sqrt{-E}x}$ to the linear Schrödinger equation (3.11) for $A > 0$ with $E < 0$ and $P(x) = 0$. Solving the differential equations (4.3) for $\sigma(t)$ and $\tau(t)$ in the case where $n = 6$, we obtain

$$\sigma(t) = \frac{1}{\sqrt{-E}} \ln\left[A\sqrt{-E}t\right] \qquad \text{and} \qquad \tau(t) = \frac{\sqrt{-E}}{2}t^2. \tag{5.1}$$

In this case the inverse of $\tau(t)$ is $f(\tau) = \sqrt{\frac{2}{\sqrt{-E}}\tau}$ and so equation (4.5) becomes $\dot{\overline{\tau}}(t) = 2\sqrt{-E}\overline{\tau}(t)$ therefore $\overline{\tau}(t) = c_0 e^{2\sqrt{-E}t}$ for any $c_0 > 0$. Using these functions and (4.6)-(4.7), we obtain

$$a(t) = \left(2c_0\sqrt{-E}\right)^{1/6} e^{\frac{1}{3}(\sqrt{-E}-\lambda-\mu)t} \tag{5.2}$$

$$b(t) = C_1\left(2c_0\sqrt{-E}\right)^{1/6} e^{\frac{1}{3}(\sqrt{-E}+2\lambda-\mu)t} \tag{5.3}$$

$$c(t) = C_2\left(2c_0\sqrt{-E}\right)^{1/6} e^{\frac{1}{3}(\sqrt{-E}-\lambda+2\mu)t} \tag{5.4}$$

for λ, μ as in (4.4). Since $P(x) = 0$, $\psi(x) = \psi_0 \equiv$ any constant therefore $\phi(t) = \psi_0$ also. Since ϕ is constant, V is also constant by (4.8) which then reduces to the formula

$$V = \frac{1}{3K^2C_1C_2}\left[(u')^2 + Eu^2\right]. \tag{5.5}$$

For this example $V = 0$, so that (5.2)-(5.4) is a vacuum solution. Note that one can show by contradiction that for any nonzero λ, μ satisfying (4.4) then $\sqrt{-E} - \lambda - \mu < 0$, $\sqrt{-E} + 2\lambda - \mu > 0$ and $\sqrt{-E} - \lambda + 2\mu > 0$.

As another example, take $u(x) = Ae^{-\sqrt{-E}x} - \frac{1}{4A\sqrt{-E}}e^{\sqrt{-E}x}$ for any real number A and any $E < 0$. Again, take $P(x) = 0$. Solving (4.3) for $n = 6$, we obtain

$$\sigma(t) = \frac{1}{2\sqrt{-E}}\ln\left[4A^2\sqrt{-E}\tanh^2\left(\frac{\sqrt[4]{-E}}{2}t\right)\right] \quad \text{and} \quad \tau(t) = \cosh(\sqrt[4]{-E}t). \tag{5.6}$$

Using that $\dot{\tau}(f(\tau)) = 1/f'(\tau) = \sqrt[4]{-E}\sqrt{\tau^2-1}$, which is obtained by differentiating the relation $f(\tau(t)) = t$ with respect to t, equation (4.5) becomes $\dot{\overline{\tau}}(t) = \sqrt{-E}(\overline{\tau}(t)^2 - 1)$ therefore $\overline{\tau}(t) = -\coth(\sqrt{-E}(t - c_0))$ for $c_0 > 0$. Using these functions and (4.6)-(4.7), we obtain

$$a(t) = (-E)^{1/12}e^{-\frac{1}{3}(\lambda+\mu)t}\operatorname{csch}^{1/3}(\sqrt{-E}(t-c_0)) \tag{5.7}$$

$$b(t) = C_1(-E)^{1/12}e^{-\frac{1}{3}(\mu-2\lambda)t}\operatorname{csch}^{1/3}(\sqrt{-E}(t-c_0)) \tag{5.8}$$

$$c(t) = C_2(-E)^{1/12}e^{-\frac{1}{3}(\lambda-2\mu)t}\operatorname{csch}^{1/3}(\sqrt{-E}(t-c_0)) \tag{5.9}$$

for $t > c_0$. Again since $P(x) = 0$, $\phi(t) = \psi_0 \equiv$ any constant and V is also a constant given by (5.5) which yields $V = \frac{\sqrt{-E}}{3K^2C_1^2C_2^2}$.

References

[1] T. Christodoulakis, T. Grammenos, C. Helias, P. Kevrekidis, G. Papadopoulos, F. Williams, Nova Science Pub. (2005).

[2] T. Christodoulakis, T. Grammenos, C. Helias, P. Kevrekidis, A. Spanou, *J. Math. Phys.*, **47** (2006) 042505.

[3] J. D'Ambroise, *Internat. J. of Pure and Applied Math.*, **42** (2008), No. 3, p. 405.

[4] J. DAmbroise, F.L. Williams, *Internat. J. of Pure and Applied Math.*, **34** (2006), No. 1, 117.

[5] R. Hawkins, J. Lidsey, *Phys. Rev.*, **D66** (2002), 023523.

[6] P. Kevrekidis, F. Williams, *Class. Quantum Gravity*, **20** (2003) L177.

[7] F. Williams, *Internat. J. of Modern Physics*, **20** (2005) 2481.

[8] F. Williams, Proceedings of 11th Marcel Grossmann Meeting, Berlin, 2006.